


### Directions for using the Olive Rectal Spray.

This product is for adults and children 12 years of age and over. For children under 12 years, ask a doctor.  
**Use before going to bed at night and if needed, repeat each morning upon waking up.**

#### Directions for rectal application

- 1 Remove seal over the bottle and cap marked "sealed for your protection".
- 2 Holding bottle upright, depress sprayer nozzle and apply a droplet on your finger to the outside of the spray tube to lubricate it prior to insertion. (The first time use of each sprayer cap may require you to pump the sprayer 3-5 times to initially fill the tube and remove the air.)
- 3 Gently insert lubricated applicator tube into the rectum as far as it will go comfortably. This can be done while lying on your back with your knees bent (as shown in the drawing) or while standing with your feet apart with your knees slightly bent. The applicator tube can be swiveled in every direction. However, this bottle is not designed to work upside-down. It is preferable to be used in the lying position so as to prevent leakage.



- 4 Suggested use: apply 2-3 sprays for each use. This sprayer will give a strong yet comfortable squirt of product. You can apply more sprays if you require. This is a homeopathic medicine and will not cause an overdose if used in higher doses than the suggested use.


- 5 REUSABLE SPRAYER - You have the option of cleaning the applicator tube after each use or replacing it with one of the 10 replacement applicators provided in the sealed pouch.

If reusing the applicator that is already on the bottle:

- be sure to clean the applicator tube after each use
- wash with soap and hot water and dry off

If you wish to replace the applicator with a new one:

- hold sprayer between fingers and gently pull upwards to disengage it from bottle. The applicator and tube will disengage from top of bottle.
  - remove a replacement applicator from the pouch and its sealed bag and gently press it onto the top of bottle.
  - do not flush the used applicator in toilet.
- Ten extra applicators are provided in the sealed pouch.



- 6 If you are able to lie down after applying the product, you can minimize leakage.

- 7 Repeat steps 1 through 6 before going to bed at night and if needed, upon waking up in the morning, for up to 7 days. If your symptoms continue beyond 7 days or do not improve after 3 days, see a doctor.

#### Directions for applying externally

- Use this product twice daily for up to 7 days as needed.
- 1 Depress pump applicator and apply a small amount of oil onto the skin outside the anus, or apply to your fingertip and then to the irritated skin, whichever way is easier for you.
  - 2 Repeat each morning and at bedtime for up to 7 days, as needed.

#### Stop use and ask your doctor if:

- symptoms do not improve in 3 days
- symptoms last more than 7 days
- you get a rash or hives, abdominal pain, fever, chills, nausea, or vomiting

These may be signs that this product is not working, or you may have a more serious condition or an allergic reaction.

If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children.** The ingredients in this product are 100% edible. However, this product is not meant to be swallowed. If swallowed, contact your doctor.

#### What side-effects may occur with Seagate Olive Rectal spray?

A mild increase in rectal burning, itching, or irritation may occur when the product is initially applied. You are applying this product to irritated and sensitive skin which tends to react when anything is initially applied.

Stop using this product and consult your doctor if you have abdominal pain, hives, skin rash, or if you have severe abdominal burning, itching, irritation, or swelling.

#### What should I do if I have questions about this product?

Questions of a medical nature, please contact your pharmacist, doctor or health care professional. If you have non-medical questions about this product or need more information on this product or any other Seagate product, call 1-888-505-4283 between 8:00 AM and 4 PM Pacific time, Monday thru Friday.

#### Other information:


- **TAMPER-EVIDENT UNIT** - Do not use if safety seal over bottle marked "sealed for your protection" is missing, has been tampered with, or if the plastic pouch containing the individually sealed replacement applicators has been opened.
- does not require refrigeration. This product can be stored or subjected to very low and high temperatures (0 F - 140 F) without affecting quality or potency. Should the product solidify when the temperature drops below 75 F follow directions above to warm the product. The natural coconut oil in this product can become solid below 75 F.
- Each spray contains 0.13 ml of product. There are 30.0 ml total product in the bottle which provides you with an approx. total of 200 sprays per bottle.



**SEAGATE®**

## OLIVE RECTAL SPRAY

- FACT SHEET -



*For the relief of anal and rectal irritations that can also be associated with yeast including:*

- Dry aching rectum
- Hemorrhoids
- Raw anus
- Pain after stool
- Swollen mucous membranes
- Burning at anus
- Soreness
- Itching
- Redness

STORE COPY

www.SeagateProducts.com  
 email: info@SeagateProducts.com

1-888-505-4283  
 5/10

### Why should I use SEAGATE Olive Rectal Spray?

Seagate Olive Rectal Spray is an all-natural homeopathic medicine designed to provide relief of the symptoms of anal and rectal irritations. This product can also be applied externally for the relief of itching and irritation on the skin outside the anal opening.

### What does Olive Rectal Spray treat?

Homeopathic medicines do not claim that they can cure or treat a disease or medical condition. Homeopathic medicines also do not claim to be antibiotic or antifungal. Homeopathic medicines can claim that they are able to offer relief for certain specific symptoms. The combination of homeopathic "active" ingredients in the Olive Rectal Spray (Aesculus hipp 12X, Aloe soc 12X, Carbo veg. 12X, Gambogia 12X and Sulphur 30X) provides relief from the symptoms of rectal irritations including:

dry aching rectum	hemorrhoids	raw anus	pain after stool
burning at anus	soreness	itching	redness
swollen mucous membranes			

Homeopathy is a natural approach to medicine that stimulates the body's immune system so that the body helps to heal itself. Under the principle of "like curing like", a homeopathic medicine will cause a mild reaction that is similar to the symptom that the body is suffering, which in turn stimulates the immune system to react. You therefore may notice immediately following the application of Olive Rectal Spray, that there will be some temporary increase in the irritating symptoms before you feel relief.

### What is the difference between Olive Rectal Spray and over-the-counter rectal and anal medications?

There are presently very few over-the-counter rectal or anal medications available to the public. Most are specifically for the temporary relief of hemorrhoid symptoms. These products come in the form of either creams, suppositories, or wipes. They contain a variety of active ingredients including: Phenylephrine HCl (a blood vessel vasoconstrictor also commonly used in sinus sprays); Hydrocortisone (a steroid hormone used to treat skin inflammation but has contraindications where there are viral or fungal diseases of the skin); Pramoxine (a "local" anesthetic used to relieve pain by deadening the nerve-endings in the skin); Petrolatum (petroleum jelly - a semi-solid mixture of hydrocarbons); and Witch Hazel (a plant-based astringent used to shrink swollen blood vessels back to their normal size).

Seagate Olive Rectal Spray uses 5 active homeopathic ingredients derived from plants: Aesculus hipp12X; Aloe soc 12X; Carbo veg. 12X; Gambogia 12X; and Sulphur 30X. Seagate combines those ingredients with these inactive ingredients: Extra Virgin Olive Oil; Coconut Oil; Omega-3 Shark Liver Oil; vanilla fragrance; and Vitamin E. All of these ingredients are natural, edible, and are completely safe even if accidentally swallowed.

### Who should first ask a doctor before using this product?

Anal and rectal irritations can occur at any age. They can be caused by some things as simple as diaper rash, constipation, hemorrhoids, or by diseases as serious as colorectal cancer. You should first ask your doctor before using this product if there is a possibility that there is a serious underlying illness or physical damage that may be causing your symptoms. Ask a doctor first before using this product if you have:

- abdominal pain and tenderness in the lower abdomen
- blood in the stool or rectal bleeding
- diarrhea, constipation, or other change in bowel habits
- intestinal obstruction
- narrow stools
- unexplained anemia
- weight loss with no known reason
- possibly ruptured your rectal membrane or wall from insertion of a foreign object
- fever chills, nausea, vomiting
- a family history of colon cancer
- been exposed to the human immunodeficiency virus (HIV) that causes AIDS

- Reoccurring anal infections (such as once a month or 3 in 6 months). You could have a serious underlying medical cause for your symptoms including cancer or a weakened immune system.

Some medical conditions can weaken the body's normal ability to fight infection. One of the most serious of these conditions is infection with human immunodeficiency virus (HIV - the virus that causes AIDS). The HIV virus causes the body to be more likely to get infections that may not clear up easily with proper treatment. If you may have been exposed to HIV, you should see your doctor right away. If you have multiple sex partners or a new sex partner, you should also ask a doctor before use to make sure you do not have an STD. For more information on HIV infection, please contact your doctor or the CDC National AIDS HOTLINE. The CDC phone numbers are: 1-800-343-AIDS (English), 1-800-344-7432 (Spanish), or 1-800-243-7889 (hearing impaired, TDD).

### What are some of the conditions that can be the cause of anal bleeding?

Gastro-intestinal bleeding, bloody stools, or bloody discharge can be caused by a variety of conditions, some that are very dangerous and some serious but not life-threatening. Dangerous conditions include: hypertension (high blood pressure leading to blood vessel rupture); bowel cancer; rectal cancer; and colon cancer. Some serious but not life-threatening conditions include: stomach ulcers; peptic ulcers; bowel inflammation; colon polyps (if benign); intestinal bleeding; rectal bleeding; prolonged constipation; and hemorrhoids. Any bleeding symptom needs professional medical advice.

### What are some conditions that have anal bleeding as a complication?

Blood discharges can also be a complication that indicates other possible underlying conditions including: Inflammatory Bowel Disease (Crohn's Disease); Cirrhosis of the Liver; Ulcers of the Large Intestine (Ulcerative colitis); Dyspepsia (inflammation of the stomach lining); Hemorrhoids; and Cancer of the Liver. Some of these conditions can be life-threatening. **Any symptom that has rectal bleeding needs professional medical advice.**

### What are the symptoms of colon or rectal infection?

Some of the symptoms of colon or rectal infection include: fever accompanying a rectal problem; a lump, sore or new growth in or around the anus; rectal bleeding without a bowel movement; or a grayish discharge. These symptoms can be the result of a virus, bacteria, or yeast.

Some drugs (such as chemotherapy or steroids) and medical conditions can weaken the body's normal ability to fight infection. One of the most serious of these conditions is with the human immunodeficiency virus (HIV- the virus that causes AIDS). Infection with HIV causes the body to be more likely to get infections, including yeast infections. If you believe you may have an infection, seek the diagnosis of your health care professional.

### Is itching a sign of infection or disease, or is the problem just a simple irritation?

Anal itching can be the result of a variety of irritations that may not be dangerous, just simply uncomfortable caused by: scented soaps or toilet paper; bleached toilet paper; poor hygiene; excessive sweating; synthetic undergarments; stress; allergic reaction to foods or vitamins; hemorrhoids; or from the reaction to other ointments. Prolonged itching, itching accompanied by pain, fever, or obstruction can also be a sign of a more serious medical condition. If you have any doubt whether the condition is just an irritation or a serious problem, you should always seek diagnosis by your health care professional.

### Can anal itching also be caused by yeast (candida)?

Rectal yeast infections can produce some or all of these symptoms: itching; burning; dryness; swelling; or abnormal discharge. Women can have some of these symptoms in the lower abdominal area or vaginal area, which may indicate either a vaginal yeast infection, a rectal yeast infection, or both. Yeast infections, (both vaginal and rectal) can be caused by: antibiotics; birth control pills; diabetes; pregnancy; sexual transmission; the reaction to soaps or douches; excessive sugar or carbohydrates; or moist undergarments.

If you believe that there is a serious cause of the itching, seek the advice of your health care professional. If you believe that you also may have a vaginal yeast infection, **do not use Olive Rectal Spray for vaginal application** even if you have changed the applicator/tube. Use Olive Rectal Spray only for rectal application. Use Seagate Olive Vaginal Spray only for Vaginal application.

### How can I get best results from using this product?

- If the temperature of the bottle drops below 75°F, the natural coconut oil can temporarily solidify in the bottle. This is normal and has no effect on quality and is the result of using only real natural raw materials in this product. Should this occur, simply warm the bottle for a few minutes (ie. in your pocket), and then gently shake the bottle before each use.
- Use product at bedtime and if needed, upon waking up in the morning, applying 2-3 sprays per application, or more if you require, for up to 7 days in a row. Symptoms should decrease with each use. Additional sprays can be applied without danger of overdose.
- Dry the genital area thoroughly after a shower, bath, or swim. Change out of a wet bathing suit or damp clothes as soon as possible. A dry area is less likely to lead to the overgrowth of yeast or bacteria.
- Wear cotton underwear and loose-fitting clothes.
- Wipe from front to back after a bowel movement or (if you are a woman) after urination .
- Do not scratch the skin outside the rectum. Scratching can cause more irritation and spread infection.
- **Tell your doctor about any drugs you are now taking.** Certain drugs such as antibiotics, steroids, and birth control pills, may make it more likely for you to get a rectal yeast infection. If you are taking any of these drugs do not stop taking them without first asking your doctor.
- If you have any other medical questions and concerns about rectal yeast infections, or the symptoms you are experiencing, call your doctor or health care professional.

### What warnings should I know about when using this product?

For rectal use only.

**Do not use if you have continual bloody discharge. Seek immediate diagnosis by a doctor.**

**Ask a doctor before use if you have:**

- **lower abdominal, back or shoulder pain, fever chills, nausea, or vomiting You may have a more serious condition.**
- recurring rectal infections (such as once a month or 3 in 6 months). You could have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system.
- been exposed to the human immunodeficiency virus (HIV) that causes AIDS

**When using this product:**

- mild increase in rectal burning, itching, or irritation may occur initially
- avoid dripping on clothing as this product will stain
- do not use if safety seal over bottle marked "sealed for your protection" is missing or has been tampered with, or if the plastic pouch containing the individually-sealed replacement applicators has been opened.